

OPERATORS THAT ATTAIN REDUCED MINIMUM

S. H. Kulkarni* and G. Ramesh**

*Department of Mathematics, Indian Institute of Technology - Palakkad,
Ahalia Integrated Campus, Kozhippara, Palakkad 678 557, India

**Department of Mathematics, Indian Institute of Technology - Hyderabad,
Kandi, Sangareddy, Telangana 502 285 India
e-mails: shk@iitpkd.ac.in; rameshg@math.iith.ac.in

(Received 15 December 2017; accepted 30 October 2019)

Let H_1, H_2 be complex Hilbert spaces and T be a densely defined closed linear operator from its domain $D(T)$, a dense subspace of H_1 , into H_2 . Let $N(T)$ denote the null space of T and $R(T)$ denote the range of T .

Recall that $C(T) := D(T) \cap N(T)^\perp$ is called the *carrier space of T* and the *reduced minimum modulus* $\gamma(T)$ of T is defined as:

$$\gamma(T) := \inf\{\|T(x)\| : x \in C(T), \|x\| = 1\}.$$

Further, we say that T attains its reduced minimum modulus if there exists $x_0 \in C(T)$ such that $\|x_0\| = 1$ and $\|T(x_0)\| = \gamma(T)$. We discuss some properties of operators that attain reduced minimum modulus. In particular, the following results are proved.

1. The operator T attains its reduced minimum modulus if and only if its Moore-Penrose inverse T^\dagger is bounded and attains its norm, that is, there exists $y_0 \in H_2$ such that $\|y_0\| = 1$ and $\|T^\dagger\| = \|T^\dagger(y_0)\|$.
2. For each $\epsilon > 0$, there exists a bounded operator S such that $\|S\| \leq \epsilon$ and $T + S$ attains its reduced minimum.

Key words : Densely defined operator; closed operator; reduced minimum modulus; minimum modulus; minimum attaining operator; reduced minimum attaining operator; gap metric; carrier graph topology; Moore-Penrose inverse.

2010 Mathematics Subject Classification : 47A05, 47A10, 47A55, 47A58.

1. INTRODUCTION

Let H_1 and H_2 be complex Hilbert spaces and $T : H_1 \rightarrow H_2$ be a bounded linear operator. We say T to be *norm attaining* if there exists $x_0 \in H_1$ such that $\|x_0\| = 1$ and $\|Tx_0\| = \|T\|$. The norm attaining operators are well studied in the literature by several authors (see [22] for details and references therein). A well known theorem in this connection is the Lindenstrauss theorem which asserts the denseness of norm attaining operators in the space of bounded linear operators between two Hilbert spaces with respect to the operator norm (see for example, [6] for a simple proof of this fact).

A natural analogue for this class of operators is the class of minimum attaining operators. Recall that a bounded operator $T : H_1 \rightarrow H_2$ is said to be *minimum attaining*, if there exists $x_0 \in H_1$ with $\|x_0\| = 1$ such that $\|Tx_0\| = m(T)$, the *minimum modulus* of T . This class of operators was first introduced by Carvajal and Neves in [5] and several basic properties were also studied in the line of norm attaining operators.

A Lindenstrauss type theorem for minimum attaining operators is proved in [16]. Moreover, rank one perturbations of closed operators is also discussed.

In this article, we define operators that attain the reduced minimum modulus and establish several basic properties of such operators. We prove that if a densely defined closed operator T attains its reduced minimum, then its Moore-Penrose inverse T^\dagger is bounded and attains its norm. It turns out that this class is a subclass of minimum attaining operators as well as the class of closed range operators. Finally, we observe that this class is dense in the class of densely defined closed operators with respect to the gap metric as well as with respect to the carrier graph topology (see [13] for details). We prove several consequences of this result.

In the second section we summarize without proofs the relevant material on densely defined closed operators, the gap metric and the carrier graph topology. In the third section we define the reduced minimum attaining operators, prove some of the basic and important properties of such operators and compare with those of minimum attaining operators. In proving most of our results, we make use of the corresponding result for minimum attaining operators, which can be found in [16] and [11].

2. PRELIMINARIES

Throughout we consider infinite dimensional complex Hilbert spaces which will be denoted by H, H_1, H_2 etc. The inner product and the induced norm are denoted by $\langle \cdot, \cdot \rangle$ and $\|\cdot\|$, respectively. The closure of a subspace M of H is denoted by \overline{M} . We denote the unit sphere of M by $S_M = \{x \in M : \|x\| = 1\}$.

Let T be a linear operator with domain $D(T)$, a subspace of H_1 and taking values in H_2 . If $D(T)$ is dense in H_1 , then T is called a *densely defined operator*.

The graph $G(T)$ of T is defined by $G(T) := \{(x, Tx) : x \in D(T)\} \subseteq H_1 \times H_2$. If $G(T)$ is closed, then T is called a *closed operator*. Equivalently, T is closed if and only if if (x_n) is a sequence in $D(T)$ such that $x_n \rightarrow x \in H_1$ and $Tx_n \rightarrow y \in H_2$, then $x \in D(T)$ and $Tx = y$.

For a densely defined operator, there exists a unique linear operator (in fact, a closed operator) $T^* : D(T^*) \rightarrow H_1$, with

$$D(T^*) := \{y \in H_2 : x \rightarrow \langle Tx, y \rangle \text{ for all } x \in D(T) \text{ is continuous}\} \subseteq H_2$$

satisfying $\langle Tx, y \rangle = \langle x, T^*y \rangle$ for all $x \in D(T)$ and $y \in D(T^*)$.

We say T to be bounded if there exists $M > 0$ such that $\|Tx\| \leq M\|x\|$ for all $x \in D(T)$. Note that if T is densely defined and bounded then T can be extended to all of H_1 in a unique way.

By the closed graph Theorem [21], an everywhere defined closed operator is bounded. Hence the domain of an unbounded closed operator is a proper subspace of a Hilbert space.

The space of all bounded linear operators between H_1 and H_2 is denoted by $\mathcal{B}(H_1, H_2)$ and the class of all densely defined, closed linear operators between H_1 and H_2 is denoted by $\mathcal{C}(H_1, H_2)$. We write $\mathcal{B}(H, H) = \mathcal{B}(H)$ and $\mathcal{C}(H, H) = \mathcal{C}(H)$.

If $T \in \mathcal{C}(H_1, H_2)$, then the null space and the range space of T are denoted by $N(T)$ and $R(T)$ respectively and the space $C(T) := D(T) \cap N(T)^\perp$ is called the *carrier* of T . In fact, $D(T) = N(T) \oplus^\perp C(T)$ [2, page 340].

Let $T_C := T|_{C(T)}$. As $\overline{C(T)} = N(T)^\perp$ (see [13, Lemma 3.3] for details), $T \in \mathcal{C}(N(T)^\perp, H_2)$.

Let $S, T \in \mathcal{C}(H)$ be operators with domains $D(S)$ and $D(T)$, respectively. Then $S + T$ is an operator with domain $D(S + T) = D(S) \cap D(T)$ defined by $(S + T)(x) = Sx + Tx$ for all $x \in D(S + T)$. The operator ST has the domain $D(ST) = \{x \in D(T) : Tx \in D(S)\}$ and is defined as $(ST)(x) = S(Tx)$ for all $x \in D(ST)$.

If S and T are closed operators with the property that $D(T) \subseteq D(S)$ and $Tx = Sx$ for all $x \in D(T)$, then T is called the *restriction* of S and S is called an *extension* of T . We denote this by $T \subseteq S$.

An operator $T \in \mathcal{C}(H)$ is said to be *normal* if $T^*T = TT^*$, *self-adjoint* if $T = T^*$, *symmetric* if $T \subseteq T^*$, *positive* if $T = T^*$ and $\langle Tx, x \rangle \geq 0$ for all $x \in D(T)$.

Let $V \in \mathcal{B}(H_1, H_2)$. Then V is called an *isometry* if $\|Vx\| = \|x\|$ for all $x \in H_1$ and a *partial isometry* if $V|_{N(V)^\perp}$ is an isometry. The space $N(V)^\perp$ is called the *initial space* or the *initial domain* and the space $R(V)$ is called the *final space* or the *final domain* of V .

If M is a closed subspace of a Hilbert space H , then P_M denotes the orthogonal projection $P_M : H \rightarrow H$ with range M .

Here we recall definition and properties of the Moore-Penrose inverse (or generalized inverse) of a densely defined closed operator that we need for our purpose.

Definition 2.1 — (Moore-Penrose Inverse) [2, Pages 314, 318-320]. Let $T \in \mathcal{C}(H_1, H_2)$. Then there exists a unique operator $T^\dagger \in \mathcal{C}(H_2, H_1)$ with domain $D(T^\dagger) = R(T) \oplus^\perp R(T)^\perp$ and has the following properties:

1. $TT^\dagger y = P_{\overline{R(T)}} y$, for all $y \in D(T^\dagger)$
2. $T^\dagger T x = P_{N(T)^\perp} x$, for all $x \in D(T)$
3. $N(T^\dagger) = R(T)^\perp$.

This unique operator T^\dagger is called the *Moore-Penrose inverse* or the *generalized inverse* of T .

The following property of T^\dagger is also well known.

For every $y \in D(T^\dagger)$, let

$$L(y) := \left\{ x \in D(T) : \|Tx - y\| \leq \|Tu - y\| \text{ for all } u \in D(T) \right\}.$$

Here any $u \in L(y)$ is called a *least square solution* of the operator equation $Tx = y$. The vector $T^\dagger y \in L(y)$, $\|T^\dagger y\| \leq \|y\|$ for all $y \in D(T^\dagger)$ and it is called the *least square solution of minimal norm*. A different treatment of T^\dagger is given in [2, Pages 336, 339, 341], where it is called “*the Maximal Tseng generalized Inverse*”.

Theorem 2.2 — [2, Page 320]. Let $T \in \mathcal{C}(H_1, H_2)$. Then

1. $D(T^\dagger) = R(T) \oplus^\perp R(T)^\perp$, $N(T^\dagger) = R(T)^\perp = N(T^*)$
2. $R(T^\dagger) = C(T)$
3. $T^\dagger \in \mathcal{C}(H_2, H_1)$
4. T^\dagger is continuous if and only if $R(T)$ is closed

5. $T^{\dagger\dagger} = T$
6. $T^{*\dagger} = T^{\dagger*}$
7. $N(T^{*\dagger}) = N(T)$
8. T^*T and $T^{\dagger}T^{*\dagger}$ are positive and $(T^*T)^{\dagger} = T^{\dagger}T^{*\dagger}$
9. TT^* and $T^{*\dagger}T^{\dagger}$ are positive and $(TT^*)^{\dagger} = T^{*\dagger}T^{\dagger}$.

Definition 2.3 — [6]. Let $T \in \mathcal{B}(H_1, H_2)$. Then T is said to be *norm attaining* if there exists $x_0 \in S_{H_1}$ such that $\|Tx_0\| = \|T\|$.

We denote the set of all norm attaining operators between H_1, H_2 by $\mathcal{N}(H_1, H_2)$ and $\mathcal{N}(H, H)$ by $\mathcal{N}(H)$.

Definition 2.4 — [2, 7, 24]. Let $T \in \mathcal{C}(H_1, H_2)$. Then

$$m(T) := \inf \{\|Tx\| : x \in S_{D(T)}\}$$

$$\gamma(T) := \inf \{\|Tx\| : x \in S_{C(T)}\},$$

are called the *minimum modulus* and the *reduced minimum modulus* of T , respectively. The operator T is said to be bounded below if and only if $m(T) > 0$.

Remark 2.5 : If $T \in \mathcal{C}(H_1, H_2)$, then

- (a) $m(T) \leq \gamma(T)$ and equality holds if T is one-to-one
- (b) $m(T) > 0$ if and only if $R(T)$ is closed and T is one-to-one.

Proposition 2.6 — [2, 10]. Let $T \in \mathcal{C}(H_1, H_2)$. Then the following statements are equivalent;

1. $R(T)$ is closed
2. $R(T^*)$ is closed
3. $T_0 := T|_{C(T)}$ has a bounded inverse
4. $\gamma(T) > 0$
5. T^{\dagger} is bounded. In fact, $\|T^{\dagger}\| = \frac{1}{\gamma(T)}$
6. $R(T^*T)$ is closed

7. $R(TT^*)$ is closed.

Remark 2.7 : If $T \in \mathcal{C}(H)$ and $T^{-1} \in \mathcal{B}(H)$, then $m(T) = \frac{1}{\|T^{-1}\|}$, by (5) of Proposition 2.6.

Theorem 2.8 — [Theorem 13.31, page 349], [3, Theorem 4, page 144]. Let $T \in \mathcal{C}(H)$ be positive. Then there exists a unique positive operator S such that $T = S^2$.

Theorem 2.9 — [3, Theorem 2, page 184]. Let $T \in \mathcal{C}(H_1, H_2)$. Then there exists a unique partial isometry $V : H_1 \rightarrow H_2$ with initial space $\overline{R(T^*)}$ and range $\overline{R(T)}$ such that $T = V|T|$.

Remark 2.10 : For $T \in \mathcal{C}(H_1, H_2)$, the operator $|T| := (T^*T)^{\frac{1}{2}}$ is called the modulus of T . Moreover, $D(|T|) = D(T)$, $N(|T|) = N(T)$ and $\overline{R(|T|)} = \overline{R(T^*)}$. As $\|Tx\| = \||T|x\|$ for all $x \in D(T)$, we can conclude that $m(T) = m(|T|)$, and $\gamma(T) = \gamma(|T|)$.

Definition 2.11 — [21, page 346]. Let $T \in \mathcal{C}(H)$. The resolvent of T is defined by

$$\rho(T) := \{\lambda \in \mathbb{C} : T - \lambda I : D(T) \rightarrow H \text{ is invertible and } (T - \lambda I)^{-1} \in \mathcal{B}(H)\}$$

and

$$\sigma(T) := \mathbb{C} \setminus \rho(T)$$

$$\sigma_p(T) := \{\lambda \in \mathbb{C} : T - \lambda I : D(T) \rightarrow H \text{ is not one-to-one}\},$$

are called the *spectrum* and the *point spectrum* of T , respectively.

The operator S is called the square root of T and is denoted by $S = T^{\frac{1}{2}}$.

Definition 2.12 — [10, Page 267]. Let $T \in \mathcal{C}(H)$. Then the *numerical range* of T is defined by

$$W(T) := \left\{ \langle Tx, x \rangle : x \in S_{D(T)} \right\}.$$

The following Proposition is proved in [17, Chapter 10] for regular (unbounded) operators between Hilbert C^* -modules, which is obviously true for densely defined closed operators in a Hilbert space.

Proposition 2.13 — [23, Lemma 5.8]. Let $T \in \mathcal{C}(H)$. Let $Q_T := (I + T^*T)^{-\frac{1}{2}}$ and $F_T := TQ_T$. Then

1. $Q_T \in \mathcal{B}(H)$ and $0 \leq Q_T \leq I$
2. $R(Q_T) = D(T)$
3. $(F_T)^* = F_T^*$

4. $\|F_T\| < 1$ if and only if $T \in \mathcal{B}(H_1, H_2)$
5. $T = F_T(I - F_T^*F_T)^{-\frac{1}{2}}$
6. $Q_T = (I - F_T^*F_T)^{\frac{1}{2}}$.

The operator F_T is called the bounded transform of T or the z -transform of T .

Lemma 2.14 — [8, 9, 19]. Let $T \in \mathcal{C}(H_1, H_2)$. Denote $\check{T} = (I + T^*T)^{-1}$ and $\hat{T} = (I + TT^*)^{-1}$. Then

1. $\check{T} \in \mathcal{B}(H_1), \hat{T} \in \mathcal{B}(H_2)$
2. $\hat{T}T \subseteq T\check{T}, \quad \|T\check{T}\| \leq \frac{1}{2}$ and $\check{T}T^* \subseteq T^*\hat{T}, \quad \|T^*\hat{T}\| \leq \frac{1}{2}$.

One of the most useful and well studied metric on $\mathcal{C}(H_1, H_2)$ is the gap metric. Here we give some details.

Definition 2.15 — (Gap between subspaces). [10, page 197]. Let H be a Hilbert space and M, N be closed subspaces of H . Let $P = P_M$ and $Q = P_N$. Then the gap between M and N is defined by

$$\theta(M, N) = \|P - Q\|.$$

If $S, T \in \mathcal{C}(H_1, H_2)$, then $G(T), G(S) \subseteq H_1 \times H_2$ are closed subspaces. The gap between $G(T)$ and $G(S)$ is called the gap between T and S . For a deeper discussion on these concepts we refer to [10, Chapter IV] and [1, page 70].

On $\mathcal{B}(H)$, the norm topology and the topology induced by the gap metric are the same. This can be seen from the following inequalities.

Theorem 2.16 — [18, Theorem 2.5]. Let $A, B \in \mathcal{B}(H)$. Then

$$\theta(A, B) \leq \|A - B\| \leq \sqrt{1 + \|A\|^2} \sqrt{1 + \|B\|^2} \theta(A, B).$$

We remark that though the above result is stated for operators defined on a Hilbert space, it remains true for operators defined between two different Hilbert spaces.

Definition 2.17 — Let $T \in \mathcal{C}(H_1, H_2)$. Define the *Carrier Graph* of T by

$$G_C(T) := \{(x, Tx) : x \in C(T)\} \subseteq H_1 \times H_2.$$

For $S, T \in \mathcal{C}(H_1, H_2)$, the gap between $G_C(S)$ and $G_C(T)$ is denoted by,

$$\eta(S, T) = \|P_{G_C(S)} - P_{G_C(T)}\|.$$

The topology induced by the metric $\eta(\cdot, \cdot)$ on $\mathcal{C}(H_1, H_2)$ is called the *Carrier Graph Topology*.

To compute $\eta(S, T)$ we can use the following formula;

Theorem 2.18 — *Let $T, S \in \mathcal{C}(H_1, H_2)$. Then*

$$|\eta(T, S) - \theta(N(T), N(S))| \leq \theta(T, S) \leq \eta(T, S) + \theta(N(T), N(S)).$$

If $N(T) = N(S)$, by Theorem 2.19, we can conclude that $\eta(S, T) = \theta(S, T)$. For the details of this metric we refer to [13].

3. MAIN RESULTS

In this section we define reduced minimum modulus attaining operators and discuss their properties. Recall that $T \in \mathcal{C}(H_1, H_2)$ is called minimum attaining if there exists $x_0 \in S_{D(T)}$ such that $\|Tx_0\| = m(T)$. In particular, if $T \in \mathcal{B}(H_1, H_2)$, then T is minimum attaining if there exists $x_0 \in S_{H_1}$ such that $\|Tx_0\| = m(T)$.

We denote the class of minimum attaining densely defined closed operators between H_1 and H_2 by $\mathcal{M}_c(H_1, H_2)$ and $\mathcal{M}_c(H, H)$ by $\mathcal{M}_c(H)$. The class of bounded minimum attaining operators is denoted by $\mathcal{M}(H_1, H_2)$ and $\mathcal{M}(H, H)$ by $\mathcal{M}(H)$.

We propose the following definition;

Definition 3.1 — We say $T \in \mathcal{C}(H_1, H_2)$ to be *reduced minimum attaining* if there exists $x_0 \in S_{C(T)}$ such that $\|Tx_0\| = \gamma(T)$.

The class of reduced minimum attaining densely defined closed linear operators between H_1 and H_2 is denoted by $\Gamma_c(H_1, H_2)$. If $H_1 = H_2 = H$, then we write $\Gamma_c(H_1, H_2)$ by $\Gamma_c(H)$. The class of bounded operators which attain the reduced minimum is denoted by $\Gamma(H_1, H_2)$ and $\Gamma(H, H)$ is denoted by $\Gamma(H)$.

Theorem 3.2 — *Let $T \in \mathcal{C}(H_1, H_2)$. Then T attains its reduced minimum if and only if T^\dagger is bounded and attains its norm.*

PROOF : Suppose T attains its reduced minimum. Then there exists $x_0 \in C(T)$ such that $\|x_0\| = 1$ and $\|T(x_0)\| = \gamma(T)$. We must have $\gamma(T) > 0$ as otherwise $x_0 \in N(T)$ will imply $x_0 = 0$, a contradiction. By Proposition 2.6, T^\dagger is bounded. Let $y_0 = T(x_0)/\|T(x_0)\| = T(x_0)/\gamma(T)$. Then $\|y_0\| = 1$ and

$$\|T^\dagger(y_0)\| = \|T^\dagger T(x_0)\|/\gamma(T) = \|x_0\|/\gamma(T) = 1/\gamma(T) = \|T^\dagger\|.$$

Thus T^\dagger attains its norm.

Conversely, assume that T^\dagger is bounded and attains its norm. Then there exists $y_0 \in H_2$ such that $\|y_0\| = 1$ and $\|T^\dagger(y_0)\| = \|T^\dagger\|$. Let $y_0 = u + v$ where $u \in R(T)$ and $v \in R(T)^\perp$. Suppose $v \neq 0$. Then $\|u\| < 1$. Hence

$$\|T^\dagger\| = \|T^\dagger(y_0)\| = \|T^\dagger(u)\| \leq \|T^\dagger\| \|u\| < \|T^\dagger\|,$$

a contradiction. This implies that $v = 0$, hence $y_0 \in R(T)$. Thus there exists $x_0 \in C(T)$ such that $y_0 = T(x_0)$. Then $x_0 = T^\dagger(y_0)$, hence $\|x_0\| = \|T^\dagger\|$. Let $z_0 = x_0/\|x_0\|$. Then $z_0 \in C(T)$, $\|z_0\| = 1$ and

$$\|T(z_0)\| = \|y_0\|/\|x_0\| = 1/\|T^\dagger\| = \gamma(T).$$

Thus T attains the reduced minimum modulus. \square

Corollary 3.3 — Let $T \in \mathcal{C}(H_1, H_2)$. Suppose T is one-to-one. Then the following are equivalent.

1. $T \in \mathcal{M}_c(H_1, H_2)$
2. $T \in \Gamma_c(H_1, H_2)$
3. $T^\dagger \in \mathcal{B}(H_2, H_1)$ and attains its norm.

PROOF : Since T is injective, $N(T) = \{0\}$. Hence $C(T) = D(T)$ and $\gamma(T) = m(T)$. This shows equivalence of (1) and (2). Equivalence of (2) and (3) follows from Theorem 3.2. \square

Lemma 3.4 — Let $T \in \mathcal{C}(H_1, H_2)$. Then $T \in \Gamma_c(H_1, H_2)$ if and only if $T_C \in \mathcal{M}_c(N(T)^\perp, H_2)$.

PROOF : The proof follows from the fact that $m(T_C) = \gamma(T)$. \square

Proposition 3.5 — Let $T \in \Gamma_c(H_1, H_2)$. Then $R(T)$ is closed.

PROOF : This follows from Theorem 3.2 and Proposition 2.6. \square

Example 3.6 : (1) All orthogonal projections on a Hilbert space attain their reduced minimum

(2) An operator with non closed range cannot attain its reduced minimum.

Proposition 3.7 — Let $T \in \mathcal{C}(H_1, H_2)$. Then $T \in \Gamma_c(H_1, H_2)$ if and only if $|T| \in \Gamma_c(H_1)$.

PROOF : By definition $D(|T|) = D(T)$ and $N(|T|) = N(T)$. Hence $C(|T|) = C(T)$. Also, $\|Tx\| = \||T|x\|$ for all $x \in D(T)$. Thus $T \in \Gamma_c(H_1, H_2)$ if and only if $|T| \in \Gamma_c(H_1)$. \square

Proposition 3.8 — [14, Proposition 4.2]. Let $T = T^* \in \mathcal{C}(H)$. Then $\gamma(T) = d(0, \sigma(T) \setminus \{0\})$.

Lemma 3.9 —

1. Let $T \in \mathcal{C}(H)$ be self-adjoint. Then $m(T) = d(0, \sigma(T))$
2. If $T \in \mathcal{C}(H_1, H_2)$, then $m(T) \in \sigma(|T|)$. In particular, if $H_1 = H_2 = H$ and $T \geq 0$, then $m(T) \in \sigma(T)$.

PROOF : Proof of (1): If T is not invertible, then $0 \in \sigma(T)$ and T is not bounded below. Hence in this case $m(T) = 0 = d(0, \sigma(T))$.

Next assume that $0 \notin \sigma(T)$. Since $\sigma(T)$ is closed ([23, Proposition 2.6, Page 29]), we can conclude that $d(0, \sigma(T)) > 0$. Also, as $T^{-1} \in \mathcal{B}(H)$, T must be bounded below. Hence $m(T) > 0$. In this case, $m(T) = \gamma(T)$. Now, by Proposition 3.8, we have $m(T) = \gamma(T) = d(0, \sigma(T) \setminus \{0\}) = d(0, \sigma(T))$.

Proof of (2): Note that $|T| \geq 0$ and by (1), we have that $m(T) = m(|T|) = d(0, \sigma(|T|))$. Since, $\sigma(|T|)$ is closed, we can conclude that $m(T) \in \sigma(|T|)$. If $H_1 = H_2 = H$ and $T \geq 0$, then we have $|T| = T$. Hence in this case the result follows. \square

Remark 3.10 : Let $T \in \mathcal{C}(H)$ be normal. Then we can prove the formula $m(T) = d(0, \sigma(T))$. First note that the crucial point in proving this in the self-adjoint case is Proposition 3.8. This is proved for normal operators in [12, Theorem 4.4.5]. Now following along the similar lines of Proposition 3.9, we can obtain the formula.

Proposition 3.11 — Let $T = T^* \in \mathcal{C}(H)$. Then $T \in \Gamma_c(H)$ if and only if either $\gamma(T)$ or $-\gamma(T)$ is an eigenvalue of T . In particular, if $T \geq 0$, then $T \in \Gamma_c(H)$ if and only if $\gamma(T)$ is an eigenvalue of T .

PROOF : We have by Lemma 3.4, that $T \in \Gamma_c(H)$ if and only if $T_C \in \mathcal{M}(N(T)^\perp)$. As $N(T)^\perp$ is a reducing subspace for T , T_C is self-adjoint. Now, $T_C \in \mathcal{M}(N(T)^\perp)$ if and only either $m(T_C)$ or $-m(T_C)$ is an eigenvalue for T_C and hence for T . Since $m(T_C) = \gamma(T)$, the conclusion follows. In particular, if $T \geq 0$, the eigenvalues of T are positive, so we can conclude that $T \in \Gamma_c(H)$ if and only if $\gamma(T) \in \sigma_p(T)$. \square

Proposition 3.12 — Let $T \in \mathcal{C}(H_1, H_2)$. Then $T \in \Gamma_c(H_1, H_2)$ if and only if $T^*T \in \Gamma_c(H_1)$.

PROOF : By Theorem 3.2, $T \in \Gamma_c(H_1, H_2)$ if and only if $R(T)$ is closed and $T^\dagger \in \mathcal{M}(H_2, H_1)$. This is equivalent to the condition that $(T^*)^\dagger = (T^\dagger)^* \in \mathcal{M}(H_1, H_2)$. This is in turn equivalent to

the fact that $(T^\dagger)(T^\dagger)^* \in \mathcal{M}(H_2)$. But $(T^\dagger)(T^\dagger)^* = (T^*T)^\dagger$, by Theorem 2.2. Thus by Theorem 3.2, $T^*T \in \Gamma_c(H_1)$. \square

Proposition 3.13 — Let $T \in \mathcal{C}(H_1, H_2)$. Then $T \in \Gamma_c(H_1, H_2)$ if and only if $T^* \in \Gamma_c(H_2, H_1)$.

PROOF : If $T \in \Gamma_c(H_1, H_2)$, then $R(T)$ is closed and so is $R(T^*)$. Also, we have $\gamma(T) = \gamma(T^*)$. By Theorem 3.2, $T^\dagger \in \mathcal{N}(H_2, H_1)$. Also, $(T^\dagger)^* \in \mathcal{N}(H_1, H_2)$, by [4, Proposition 2.5]. Note that $(T^\dagger)^* = (T^*)^\dagger$, by Theorem 2.2. Hence by Theorem 3.2 again, $T^* \in \Gamma_c(H_2, H_1)$. Applying the same result for T^* and observing that $T^{**} = T$, we get the other way implication. \square

Remark 3.14 : The above result need not hold for minimum attaining operators. Let $H = \ell^2$ and $\{e_n : n \in \mathbb{N}\}$ denote the standard orthonormal basis for H . That is $e_n(m) = \delta_{nm}$, the Kronecker delta function. Define operators $D, R : H \rightarrow H$ by

$$De_n = \frac{1}{n}e_n,$$

$$Re_n = e_{n+1} \text{ for each } n \in \mathbb{N}.$$

Let $T = RD$. Since R is an isometry, we have $m(T) = m(D) = d(0, \sigma(D)) = \inf \left\{ \frac{1}{n} : n \in \mathbb{N} \right\} = 0$. Since $0 \notin \sigma_p(D)$, D is not minimum attaining. Thus T is not minimum attaining. But, $N(T^*) = \text{span}\{e_1\}$. So $m(T^*) = 0$ and $T^* \in \mathcal{M}(H)$. Note that $T^*T = D^2$ cannot have closed range since D is compact. Equivalently, $R(T)$ is not closed, whence T cannot attain its reduced minimum by Theorem 3.2.

Proposition 3.15 — Let $T \in \mathcal{C}(H_1, H_2)$. If $T \in \Gamma_c(H_1, H_2)$, then $T \in \mathcal{M}_c(H_1, H_2)$.

PROOF : First assume that T is one-to-one. Then $\gamma(T) = m(T)$. Hence if $T \in \Gamma_c(H_1, H_2)$, then clearly $T \in \mathcal{M}_c(H_1, H_2)$. If T is not one-to-one, then $N(T) \neq \{0\}$. Hence in this case $m(T) = 0$ and there exists $0 \neq x \in N(T)$ such that $Tx = 0$. Hence clearly $T \in \mathcal{M}_c(H_1, H_2)$. This completes the proof. \square

Proposition 3.16 — [11, Proposition 3.5]. Let $T \in \mathcal{C}(H)$ be positive. Then

$$m(T) = \inf \{ \langle Tx, x \rangle : x \in S_{D(T)} \}.$$

In particular, if $T \in \mathcal{C}(H_1, H_2)$, then $m(T^*T) = m(T)^2$.

Proposition 3.17 — Let $T \in \mathcal{C}(H)$ be positive. Then

$$\gamma(T) = \inf \{ \langle Tx, x \rangle : x \in S_{C(T)} \}.$$

In particular, if $T \in \mathcal{C}(H_1, H_2)$, then $\gamma(T^*T) = \gamma(T)^2$.

PROOF : Since, T_C is positive, we have by Proposition 3.16,

$$\begin{aligned}\gamma(T) &= m(T_C) = \inf \{ \langle T_C x, x \rangle : x \in S_{C(T)} \} \\ &= \inf \{ \langle T x, x \rangle : x \in S_{C(T)} \}.\end{aligned}$$

Further, if $T \in \mathcal{C}(H_1, H_2)$, then $T^*T \in \mathcal{C}(H_1)$ is positive. Thus by applying the above formula for T^*T and by the definition of $\gamma(T)$, we get the conclusion.

Remark 3.18 : Let $T \in \mathcal{C}(H)$ be positive. Then the following statements are equivalent (see [11, Proposition 3.8]):

- (1) $T \in \mathcal{M}_c(H)$
- (2) $m(T)$ is an eigenvalue of T
- (3) $m(T)$ is an extreme point of $W(T)$.

In general if $T \in \Gamma_c(H)$ and is positive, then $\gamma(T)$ need not be an extreme point of the numerical range of T . To see this, consider the operator T on \mathbb{C}^3 , whose matrix with respect to the standard orthonormal basis of \mathbb{C}^3 is $\begin{pmatrix} 0 & 0 & 0 \\ 0 & \frac{1}{2} & 0 \\ 0 & 0 & 1 \end{pmatrix}$. It can be easily computed that $\gamma(T) = \frac{1}{2}$, which is not an extreme point of $W(T) = [0, 1]$, but $m(T) = 0$, which is an extreme point of $W(T)$.

Proposition 3.19 — Let $T \in \mathcal{C}(H_1, H_2)$ and F_T be the bounded transform of T . Then

- (1) $\gamma(F_T) = \frac{\gamma(T)}{\sqrt{1 + \gamma(T)^2}}$
- (2) $m(F_T) = \frac{m(T)}{\sqrt{1 + m(T)^2}}$.

PROOF : Proof of (1): In view of Proposition 3.17 it is enough to show that $\gamma(F_T^*F_T) = \frac{\gamma(T^*T)}{1 + \gamma(T^*T)}$. First we note that $F_T^*F_T = T^*T(I + T^*T)^{-1} = I - (I + T^*T)^{-1}$. Using the formula in Proposition 3.8, we get

$$\begin{aligned}\gamma(F_T^*F_T) &= d(0, \sigma(F_T^*F_T) \setminus \{0\}) \\ &= \inf \left\{ \frac{\mu}{1 + \mu} : \mu \in \sigma(T^*T) \setminus \{0\} \right\} \\ &= \inf \left\{ 1 - \frac{1}{1 + \mu} : \mu \in \sigma(T^*T) \setminus \{0\} \right\}\end{aligned}$$

$$\begin{aligned}
 &= 1 - \sup \left\{ \frac{1}{1 + \mu} : \mu \in \sigma(T^*T) \setminus \{0\} \right\} \\
 &= 1 - \frac{1}{1 + \inf \{ \mu : \mu \in \sigma(T^*T) \setminus \{0\} \}} \\
 &= \frac{\gamma(T^*T)}{1 + \gamma(T^*T)}.
 \end{aligned}$$

Hence we can conclude that $\gamma(F_T) = \frac{\gamma(T)}{\sqrt{1 + \gamma(T)^2}}$.

Proof of (2): To prove this we need to use (1) of Lemma 3.9 and follow the similar steps as above. \square

Proposition 3.20 — Let $T \in \mathcal{C}(H_1, H_2)$. Then $T \in \Gamma_c(H_1, H_2)$ if and only if $F_T \in \Gamma(H_1, H_2)$.

PROOF : In view of Proposition 3.12, it suffices to show that $T^*T \in \Gamma_c(H_1)$ if and only if $F_T^*F_T \in \Gamma(H_1)$. First, note that $(F_T)^* = F_{T^*}$. If $T^*T \in \Gamma_c(H_1)$, there exists $x_0 \in S_{\mathcal{C}(T^*T)}$ such that $T^*Tx_0 = \gamma(T^*T)x_0$. Then we have

$$\begin{aligned}
 F_T^*F_Tx_0 &= (I - (I + T^*T)^{-1})x_0 \\
 &= \frac{\gamma(T^*T)}{1 + \gamma(T^*T)}x_0 \\
 &= \frac{\gamma(T)^2}{1 + \gamma(T)^2}x_0 \\
 &= \gamma(F_T)^2x_0 \\
 &= \gamma(F_T^*F_T)x_0.
 \end{aligned}$$

This shows that $F_T^*F_T \in \Gamma(H_1)$.

To prove the converse, suppose $F_T^*F_T \in \Gamma(H_1)$. Then $F_T^*F_T = F_{T^*}F_T = T^*T(I + T^*T)^{-1} \in \Gamma(H_1)$. Thus there exists $x_0 \in N((F_T)^*F_T)^\perp = N(F_T)^\perp = N(T)^\perp$ such that $T^*T(I + T^*T)^{-1}x_0 = \gamma(F_T^*F_T)x_0$. By Proposition 3.19, we can obtain that

$$(I - (I + T^*T)^{-1})(x_0) = \left(1 - \frac{1}{1 + \gamma(T)^2}\right)(x_0).$$

Equivalently, $(I + T^*T)^{-1}(x_0) = \frac{1}{1 + \gamma(T)^2}x_0$. That is $x_0 \in R((I + T^*T)^{-1}) = D(I + T^*T) = D(T^*T)$. It follows that $(I + T^*T)(x_0) = (1 + \gamma(T)^2)(x_0)$ or $T^*Tx_0 = \gamma(T^*T)x_0$, concluding T^*T attains its reduced minimum and so is T . \square

Next, we would like to prove a Lindenstrauss type theorem for the class of reduced minimum attaining operators. We need the following results for this purpose.

Theorem 3.21 — [16, Theorem 3.1], [20, Remark 3.7]. Let $S, T \in \mathcal{C}(H_1, H_2)$ and $D(S) = D(T)$. Then

1. the operators $\widehat{T}^{\frac{1}{2}}(T - S)\check{S}^{\frac{1}{2}}$ and $\widehat{S}^{\frac{1}{2}}(T - S)\check{T}^{\frac{1}{2}}$ are bounded and

$$\theta(S, T) = \max \left\{ \|\widehat{T}^{\frac{1}{2}}(T - S)\check{S}^{\frac{1}{2}}\|, \|\widehat{S}^{\frac{1}{2}}(T - S)\check{T}^{\frac{1}{2}}\| \right\}$$

2. if $T - S$ is bounded, then $\theta(S, T) \leq \|S - T\|$.

Theorem 3.22 — [16, Theorem 3.5]. Let $T \in \mathcal{C}(H_1, H_2)$. Then for each $\epsilon > 0$, there exists $S \in \mathcal{B}(H_1, H_2)$ with $\|S\| \leq \epsilon$ such that $S + T$ is minimum attaining and $\theta(S + T, T) \leq \epsilon$. Moreover, if $m(T) > 0$, then we can choose S to be a rank one operator.

Theorem 3.23 — Let $T \in \mathcal{C}(H_1, H_2)$. Then for each $\epsilon > 0$ there exists $S \in \mathcal{B}(H_1, H_2)$ such that

1. $\|S\| \leq \epsilon$
2. $N(T) = N(T + S)$ and
3. $T + S$ attains reduced minimum, and $\theta(S + T, T) \leq \epsilon$

Moreover, if $\gamma(T) > 0$, then we can choose S to be a rank one operator.

PROOF : First assume that $\gamma(T) > 0$. Consider $T_C := T|_{C(T)} : N(T)^\perp \rightarrow H_2$ is densely defined closed operator. We may assume that $0 < \epsilon < \gamma(T)$. By Theorem 3.22, there exists $S_0 \in \mathcal{B}(N(T)^\perp, H_2)$ such that $\|S_0\| \leq \epsilon$ and $T_C + S_0$ is minimum attaining. That is, there exists $x_0 \in D(T_C + S_0) = D(T_C) = C(T)$ such that $\|x_0\| = 1$ and $\|(T_C + S_0)(x_0)\| = m(T_C + S_0)$. As $m(T_C) = \gamma(T) > 0$, we can choose S_0 to be a rank one operator.

For $x = u + v \in H_1$ with $u \in N(T)$, $v \in N(T)^\perp$, define $Sx = S_0v$. Then $\|Sx\| = \|S_0v\| \leq \|S_0\|\|v\| \leq \epsilon\|x\|$. Thus $\|S\| \leq \epsilon$. Note that S is a rank one operator.

We claim that $T + S$ attains reduced minimum. Note that $D(T + S) = D(T)$. Let $u \in N(T)$. Then $(T + S)(u) = Tu + Su = 0$. Thus $N(T) \subseteq N(T + S) = \{x \in D(T) : Tx + Sx = 0\}$. Suppose that $x \in D(T) \setminus N(T)$. Let $x = u + v$ with $u \in N(T)$ and $v \in N(T)^\perp$. Then $v \neq 0$. Also, $v \in C(T)$ as $x, u \in D(T)$. Then

$$\begin{aligned} \|(T + S)(x)\| &= \|Tv + Sv\| \\ &= \|Tv + S_0v\| \\ &\geq \|Tv\| - \|S_0v\| \end{aligned}$$

$$\begin{aligned}
 &\geq \gamma(T)\|v\| - \|S_0\|\|v\| \\
 &\geq (\gamma(T) - \epsilon)\|v\| \\
 &> 0.
 \end{aligned}$$

Thus $(T + S)(x) \neq 0$. Thus $x \notin N(T + S)$. This show that $N(T) = N(T + S)$. Since $D(T + S) = D(T)$, we have $C(T + S) = C(T)$ and hence

$$\begin{aligned}
 \gamma(T + S) &= \inf \{ \|(T + S)(x)\| : x \in C(T), \|x\| = 1 \} \\
 &= \inf \{ \|(T_C + S_0)(x)\| : x \in C(T), \|x\| = 1 \} \\
 &= \|(T_C + S_0)(x_0)\| \\
 &= \|(T + S)(x_0)\|.
 \end{aligned}$$

Next suppose that $\gamma(T) = 0$. Let $\epsilon > 0$. Choose $x_0 \in C(T)$ such that $\|x_0\| = 1$ and $\|Tx_0\| < \frac{\epsilon}{4}$. Then

$$\|(T + \frac{\epsilon}{2}I)(x_0)\| \geq \frac{\epsilon}{2} - \|Tx_0\| \geq \frac{\epsilon}{4}.$$

Hence

$$0 < \frac{\epsilon}{4} \leq m(T + \frac{\epsilon}{2}I) \leq \gamma(T + \frac{\epsilon}{2}).$$

By above argument, there exists $\tilde{S} \in \mathcal{B}(H_1, H_2)$ such that $\|\tilde{S}\| \leq \frac{\epsilon}{2}$ and $T + \frac{\epsilon}{2}I + \tilde{S}$ attains reduced minimum. Then $\|\frac{\epsilon}{2}I + \tilde{S}\| \leq \epsilon$. Take $S = \frac{\epsilon}{2}I + \tilde{S}$. Then by Theorem 3.21, S satisfies all the stated conditions. \square

We have the following consequences.

Theorem 3.24 — *The following statements holds true;*

1. $\Gamma_c(H_1, H_2)$ is dense in $\mathcal{C}(H_1, H_2)$ with respect to the gap metric $\theta(\cdot, \cdot)$
2. $\Gamma_c(H_1, H_2)$ is dense in $\mathcal{C}(H_1, H_2)$ with respect to the metric $\eta(\cdot, \cdot)$
3. the set of all closed range operators of $\mathcal{C}(H_1, H_2)$ is dense in $\mathcal{C}(H_1, H_2)$ with respect to the metric $\theta(\cdot, \cdot)$
4. the set of all closed range operators of $\mathcal{C}(H_1, H_2)$ is dense in $\mathcal{C}(H_1, H_2)$ with respect to the metric $\eta(\cdot, \cdot)$.

PROOF : Proof of (1): Follows by Theorem 3.23.

Proof of (2): Let $\epsilon > 0$. Then by Theorem 3.23, we can obtain $S \in \mathcal{B}(H_1, H_2)$ with $\|S\| \leq \epsilon$ such that $N(T) = N(T + S)$ and $\theta(T, T + S) \leq \epsilon$. By Theorem 2.19, it follows that $\eta(T + S, T) = \theta(T + S, T) \leq \epsilon$. Hence the claim.

Proof of (3): Since $\Gamma_c(H_1, H_2)$ is a subset of the set of all closed range operators in $\mathcal{C}(H_1, H_2)$, the conclusion is immediate by (2) above. \square

Proof of (4): This follows by Proposition 3.5 and (2) above. \square

Using the equivalence of the gap metric and the metric induced by the operator norm on $\mathcal{B}(H_1, H_2)$ we can obtain the following consequences.

Corollary 3.25 — The following statements are true.

1. $\Gamma(H_1, H_2)$ is dense in $\mathcal{B}(H_1, H_2)$ with respect to the operator norm
2. the set of all bounded closed range operators is dense in $\mathcal{B}(H_1, H_2)$ with respect to the operator norm.

REFERENCES

1. N. I. Akhiezer and I. M. Glazman, *Theory of linear operators in Hilbert space*, Translated from the Russian and with a preface by Merlynd Nestell. Reprint of the 1961 and 1963 translations. Two volumes bound as one. Dover Publications, Inc., New York, 1993.
2. A. Ben-Israel and T. N. E. Greville, *Generalized inverses: theory and applications*, Wiley-Interscience, New York, 1974.
3. M. Š. Birman and M. Z. Solomjak, *Spectral theory of selfadjoint operators in Hilbert space* (Russian), Leningrad. Univ., Leningrad, 1980.
4. Carvajal, Xavier; Neves, Wladimir. Operators that achieve the norm, *Integral Equations Operator Theory*, **72**(2) (2012), 179-195.
5. X. Carvajal and W. Neves, Operators that attain their minima, *Bull. Braz. Math. Soc. (N.S.)*, **45**(2) (2014), 293-312.
6. P. Enflo, J. Kover, and L. Smithies, Denseness for norm attaining operator-valued functions, *Linear Algebra Appl.*, **338** (2001), 139-144.
7. S. Goldberg, *Unbounded linear operators: Theory and applications*, McGraw-Hill, New York, 1966.
8. S. Gramsch and E. Schock, Ill-posed equations with transformed argument, *Abstr. Appl. Anal.*, **2003**(13) 785-791.

9. C. W. Groetsch, Spectral methods for linear inverse problems with unbounded operators, *J. Approx. Theory*, **70**(1) (1992), 16-28.
10. T. Kato, *Perturbation theory for linear operators*, second edition, Springer, Berlin, 1976.
11. S. H. Kulkarni and G. Ramesh, On absolutely minimum attaining closed operators, *J. Anal.*, (2019); <https://doi.org/10.1007//341478-019-00189-x>.
12. G. Ramesh, *Approximation methods for solving operator equations involving unbounded operators*, Ph. D Thesis, IIT Madras 2008.
13. S. H. Kulkarni and G. Ramesh, The carrier graph topology, *Banach J. Math. Anal.*, **5**(1) (2011), 56-69.
14. S. H. Kulkarni, M. T. Nair, and G. Ramesh, Some properties of unbounded operators with closed range, *Proc. Indian Acad. Sci. Math. Sci.*, **118**(4) (2008), 613-625.
15. S. H. Kulkarni and G. Ramesh, A formula for gap between two closed operators, *Linear Algebra Appl.*, **432**(11) (2010), 3012-3017.
16. S. H. Kulkarni and G. Ramesh, On the denseness of minimum attaining operators, *Oper. Matrices*, **12**(3) (2018), 699-709.
17. E. C. Lance, *Hilbert C^* -modules*, London Mathematical Society Lecture Note Series, 210, Cambridge Univ. Press, Cambridge, 1995.
18. R. Nakamoto, Gap formulas of operators and their applications, *Math. Japon.*, **42**(2) (1995), 219-232.
19. G. K. Pedersen, *Analysis now*, Graduate Texts in Mathematics, 118, Springer, New York, 1989.
20. G. Ramesh, McIntosh formula for the gap between regular operators, *Banach J. Math. Anal.*, **7**(1) (2013), 97-106.
21. W. Rudin, *Functional analysis*, second edition, International Series in Pure and Applied Mathematics, McGraw-Hill, New York, 1991.
22. S. Shkarin, Norm attaining operators and pseudospectrum, *Integral Equations Operator Theory*, **64**(1) (2009), 115-136.
23. Schmüdgen, Konrad, Unbounded self-adjoint operators on Hilbert space, Graduate Texts in Mathematics, 265, Springer, Dordrecht, 2012.
24. A. E. Taylor, The minimum modulus of a linear operator, and its use for estimates in spectral theory, *Studia Math. (Ser. Specjalna) Zeszyt*, **1** (1963), 131-132.